

### 35. CLEBSCH-GORDAN COEFFICIENTS, SPHERICAL HARMONICS, AND $d$ FUNCTIONS

Note: A square-root sign is to be understood over every coefficient, e.g., for  $-8/15$  read  $-\sqrt{8/15}$ .

Notation:

$J$	$J$	$\dots$
$M$	$M$	$\dots$

  

$m_1$	$m_2$		
$m_1$	$m_2$	Coefficients	
$\vdots$	$\vdots$		
$\vdots$	$\vdots$		

$Y_1^0 = \sqrt{\frac{3}{4\pi}} \cos\theta$

$Y_1^1 = -\sqrt{\frac{3}{8\pi}} \sin\theta e^{i\phi}$

$Y_2^0 = \sqrt{\frac{5}{4\pi}} \left(\frac{3}{2} \cos^2\theta - \frac{1}{2}\right)$

$Y_2^1 = -\sqrt{\frac{15}{8\pi}} \sin\theta \cos\theta e^{i\phi}$

$Y_2^2 = \frac{1}{4} \sqrt{\frac{15}{2\pi}} \sin^2\theta e^{2i\phi}$

$Y_\ell^{-m} = (-1)^m Y_\ell^{m*}$

$d_{m,0}^\ell = \sqrt{\frac{4\pi}{2\ell+1}} Y_\ell^m e^{-im\phi}$

$\langle j_1 j_2 m_1 m_2 | j_1 j_2 JM \rangle$   
 $= (-1)^{J-j_1-j_2} \langle j_2 j_1 m_2 m_1 | j_2 j_1 JM \rangle$

  

$d_{m',m}^j = (-1)^{m-m'} d_{m,m'}^j = d_{-m,-m'}^j$

$d_{0,0}^1 = \cos\theta$

$d_{1/2,1/2}^{1/2} = \cos\frac{\theta}{2}$

$d_{1,1}^1 = \frac{1+\cos\theta}{2}$

$d_{1/2,-1/2}^{1/2} = -\sin\frac{\theta}{2}$

$d_{1,0}^1 = -\frac{\sin\theta}{\sqrt{2}}$

$d_{1,-1}^1 = \frac{1-\cos\theta}{2}$

  

$d_{3/2,3/2}^{3/2} = \frac{1+\cos\theta}{2} \cos\frac{\theta}{2}$

$d_{3/2,1/2}^{3/2} = -\sqrt{3} \frac{1+\cos\theta}{2} \sin\frac{\theta}{2}$

$d_{3/2,-1/2}^{3/2} = \sqrt{3} \frac{1-\cos\theta}{2} \cos\frac{\theta}{2}$

$d_{3/2,-3/2}^{3/2} = -\frac{1-\cos\theta}{2} \sin\frac{\theta}{2}$

$d_{1/2,1/2}^{3/2} = \frac{3\cos\theta-1}{2} \cos\frac{\theta}{2}$

$d_{1/2,-1/2}^{3/2} = -\frac{3\cos\theta+1}{2} \sin\frac{\theta}{2}$

$d_{2,2}^2 = \left(\frac{1+\cos\theta}{2}\right)^2$

$d_{2,1}^2 = -\frac{1+\cos\theta}{2} \sin\theta$

$d_{2,0}^2 = \frac{\sqrt{6}}{4} \sin^2\theta$

$d_{2,-1}^2 = -\frac{1-\cos\theta}{2} \sin\theta$

$d_{2,-2}^2 = \left(\frac{1-\cos\theta}{2}\right)^2$

$d_{1,1}^2 = \frac{1+\cos\theta}{2} (2\cos\theta - 1)$

$d_{1,0}^2 = -\sqrt{\frac{3}{2}} \sin\theta \cos\theta$

$d_{1,-1}^2 = \frac{1-\cos\theta}{2} (2\cos\theta + 1)$

$d_{0,0}^2 = \left(\frac{3}{2} \cos^2\theta - \frac{1}{2}\right)$

**Figure 35.1:** The sign convention is that of Wigner (*Group Theory*, Academic Press, New York, 1959), also used by Condon and Shortley (*The Theory of Atomic Spectra*, Cambridge Univ. Press, New York, 1953), Rose (*Elementary Theory of Angular Momentum*, Wiley, New York, 1957), and Cohen (*Tables of the Clebsch-Gordan Coefficients*, North American Rockwell Science Center, Thousand Oaks, Calif., 1974). The coefficients here have been calculated using computer programs written independently by Cohen and at LBNL.